

BABALA: Ang produktong ito ay maaaring makalason kapag nalunok. Huwag kumain, uminom o manigarilyo habang nagbobomba. Iwasang malanghap, malagyan ang mga mata at madikit sa balat ang kemikal. Magsuot ng damit pananggalang habang ginagamit ito. Hugasan ng sabon at tubig ang kamay, braso at mukha bago kumain o manigarilyo. Huwag gamitin uli ang pinagsidlan ng produkto. Iwasang maapektuhan ng kemikal at ng mga sisidlan nito ang anumang pinagkukunan ng tubig.

FIRST AID:

Remove contaminated clothing.

If inhaled: Keep patient calm, move to fresh air, seek medical attention.

On skin contact: Wash thoroughly with soap and water

On contact with eyes: Wash affected eyes for at least 15 minutes under running water with eyelids held open.

On ingestion: Immediately rinse mouth and then drink 200-300 ml of water, seek medical attention.

NOTE TO PHYSICIAN: Treat according to symptoms (decontamination, vital functions).

No known specific antidote.

IN CASE OF EMERGENCY, PLEASE CONTACT:

BASF Emergency Contact Details

National Emergency No.:

+632 8831 5576

International Emergency No.:

+49 180 2273-112

National Poison Management and Control Center

PGH, Manila

Tel: (02) 8554 8400 loc. 3976,

(02) 8524 1078

Globe: +63 966 718 9904

DIMETHOMORPH
AMETOCTRADIN

GROUP 40 & 45 FUNGICIDE

BASF

We create chemistry

Zampro[®] DM 52.5SC
Fungicide

Active ingredients:
Ametoctradin..... 300 g/L
Dimethomorph..... 225 g/L
Inert ingredients..... 139 g/L
Water (as solvent)..... up to 1 L

REGISTERED WITH THE FERTILIZER AND
PESTICIDE AUTHORITY PURSUANT TO P.D. 1144

FPA REGISTRATION NUMBER: 005-425-2948

Net Contents:
250mL

KEEP OUT OF REACH OF CHILDREN



GENERAL INFORMATION: Zampro[®] DM is an innovative fungicide with a new chemical class which is highly active for the control of comycetes (like downy mildew and late blight diseases) and with excellent selectivity. Zampro[®] DM is a combination of Ametoctradin and Dimethomorph active ingredients. This product should be applied as a preventive control.

DIRECTIONS FOR USE:

CROP	DISEASE	PRODUCT RATE		APPLICATION INTERVAL
		(mL/Ha)	TBSP* /16L	
Potato	Late Blight	1000-1200	8-10**	7-10 days

* 1TBSP = 10mL

**at 200 L/ha Spray Volume

MIXING: Mix the required amount of Zampro[®] DM product into a knapsack sprayer with 16 Liter water capacity. Zampro[®] DM can be applied 6 to 8 times per season. **MAXIMUM NO. OF SEQUENTIAL SPRAY APPLICATION:** 2 times **NUMBER OF APPLICATION:** 3 times per cropping season **APPLICATION INTERVAL:** 7 days **COMPATIBILITY:** Zampro[®] DM is compatible with other fungicides. **PRE-HARVEST INTERVAL:** 14 days **RE-ENTRY:** After drying of spray droplets up to 24 hours. **STORAGE AND DISPOSAL:** Store in original container, tightly closed, and keep away from food, drinks and feeds by storage or disposal. Keep out of reach of children. Triple rinse container and dispose of in accordance with national and local regulations and requirements. **PROHIBITION AND WARRANTY:** It is a violation of FPA rules and regulations to use this product in a manner inconsistent with its label. No warranty of any kind, express or implied, is made concerning the use of this product. The user assumes all risks and liabilities resulting from handling, use or application.

Zampro[®] is a registered trademark of BASF

Registered by:
BASF Philippines, Inc.

Units 3 & 4, Upper Penthouse, CTP ASEAN Tower Blk 3 Lot 2, Spectrum District, Filinvest Corporate City Alabang, Muntinlupa City, 1781 Metro Manila, Philippines

SCAN ME!



Lot/Batch No.:

Date Formulated:

Expiry Date:

