



**BASF**

The Chemical Company

# Cumora

## 50 SC

### A SYSTEMIC FUNGICIDE FOR THE CONTROL OF BLACK SIGATOKA IN BANANA



**ACTIVE INGREDIENT:**  
Boscalid  
Inert Ingredient

500 g/L  
500 g/L

NET CONTENT:

# 20 Liters



Registered with the Fertilizer and Pesticide Authority Pursuant to P.D. 1144  
FPA Registration Number: 005-413-2598

Cumora™ is a trademark of BASF

81055175 PH 1011

### KEEP OUT OF REACH OF CHILDREN





**Precautions:** Do not get in eyes, on skin or on clothing. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing the spray mist. Do not eat, drink or smoke while working. Wear protective clothing such as long-sleeved shirts and pants when using this product. Wash with soap and water after working.

**Babala:** Ang produktong ito ay maaaring makalason kapag nalunok, nalanghap, o napunta sa balat. Huwag kumain uminom o manigarilyo habang nagbobomba. Iwasang malanghap, malagyan ang mga mata at madikit sa balat ang kemikal. Magsuot ng damit pananggalang habang ginagamit ito. Hugasan ng sabon at tubig ang kamay, braso at mukha bago kumain o manigarilyo. Huwag gamitin muli ang pinagsidlan ng produkto. Iwasang maapektuhan ng kemikal at ng mga sisidlan nito ang anumang pinagkukunan ng tubig.

**First Aid:** Remove contaminated clothing immediately. *If inhaled*, keep patient calm, bring to fresh air. Get medical attention. *On contact with eyes*, immediately wash affected eyes for 15 minutes with running water with eyelids held open. Consult an eye specialist. Get medical attention immediately. *On contact with skin*, immediately wash with soap and water. Get medical attention if irritation persists. *On ingestion*, immediately rinse mouth and drink plenty of water. Do not induce vomiting. Never give anything through mouth to an unconscious person. Call a physician or a poison control center immediately.

**IN CASE OF EMERGENCY, PLEASE CONTACT:**

<p><b>BASF Philippines, Inc.</b> 11/F Hanjinphil Corporation Building 1128 University Parkway, North Bonifacio, Global City, Taguig, Metro Manila 1634 Tel.: (02) 811 8000; Fax: (02) 838 1025</p>	<p><b>National Poison Management and Control Center</b> PGH, Manila Tel.: (02) 554 1078; (02) 521 8400 loc 2311</p>
<p><b>Vicente Sotto Medical Center</b> B. Rodriguez St., Cebu City Tel.: (032) 253 9891 loc 353</p>	<p><b>Davao Medical Center</b> Bajada, Davao City Tel.: (082) 227 2731</p>

**Note to Physician:** Treat according to symptoms (decontamination, vital functions), no known specific antidote.

**Storage and Disposal:** Store in original container, tightly closed, away from food and feeds. Keep out of reach of children. Do not contaminate water, food or feed by storage or disposal. Triple rinse container. Then offer for recycling or puncture and dispose of in a sanitary landfill.

**Prohibition and Warranty:** IT IS A VIOLATION OF FPA RULES AND REGULATIONS TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABEL. No warranty of any kind, expressed or implied is made concerning the use of this product. User assumes all risks and liabilities resulting from handling, use or application.

**Cumora® 50 SC** is a systemic fungicide which resembles the activity of strobilurin group containing 500 g/L Boscalid. It has preventive, curative and eradication action effective in controlling Black Sigatoka in banana.

**Directions for Use:** **Cumora® 50 SC** should only be applied in a spray program with other fungicides (e.g. Calixin 86 OL and Opal 7.5 EC) use in banana to control Black Sigatoka.

CROP	DISEASE	DOSAGE RATE (L/HA)	APPLICATION INTERVAL
Banana	Sigatoka	0.5 - 0.7	10-14 days

**Mixing:** Recommended mixing procedures for **Cumora® 50 SC** are as follows: 1) Add banana oil plus recommended rate of emulsifier then agitate; 2) Add 50% of total water then agitate; 3) Add the recommended rate of Cumora® 50 SC then agitate; and 4) Add the rest of the water to complete the total volume.

**Compatibility:** **Cumora® 50 SC** is compatible with banana oil (e.g. Banole) and other commonly used fungicides in banana e.g Opal and Calixin.

**PRE-HARVEST INTERVAL:** 0 days.

**RE-ENTRY PERIOD:** After drying of spray or 24 hours after spraying.

**FOR INSTITUTIONAL USE ONLY**

Cumora™ 50 SC is a trademark of BASF

Registered by:

BASF Philippines, Inc.  
11/F Hanjinphil Corporation Building  
1128 University Parkway, North Bonifacio  
Global City, Taguig, Metro Manila 1634  
Tel.: (02) 811 8000 / Fax: (02) 838 1025



Lot/Batch No:

Formulation Date:

58641418

81055186 PH 1011

