

PRECAUTIONS: Wear protective clothing particularly gloves when mixing or applying this product. Avoid skin and eye contamination, or inhalation of spray mist. Wash hands and exposed skin before meal and after work. Avoid contamination of any water supply with chemical or empty container. Very toxic to aquatic organisms.

KAUKULANG PAG-IINGAT: Magsuot ng damit pananggalang kapag naghahalo o nagbobomba. Huwag kumain, uminom o manigarilyo habang gumagamit nito. Iwasang madikit sa balat at mata o malanghap. Maglinis ng katawan bago kumain at pagkatapos magbomba. Iwasang malagyan ng kemikal ang anumang pinagkukunan ng tubig.

FIRST AID

Ingestion: If patient is conscious, give him one glass of water and induce vomiting by pressing back of tongue with a blunt instrument.


Inhalation: Remove from contaminated area and bring patient to fresh air and cover him well.

Skin Contact: Remove contaminated clothing and wash affected area with soap and water.

Eye contamination: Flush eyes with plenty of clean running water for at least 15 minutes.


Get medical attention.

IN CASE OF EMERGENCY, PLEASE CONTACT:			
BASF PHILIPPINES INC.	NATIONAL POISON AND MANAGEMENT CONTROL CENTER	VICENTE SOTTO MEDICAL CENTER	DAVAO MEDICAL CENTER
11/F HHIC BUILDING University Parkway, North Bonifacio, Global City, 1634 Taguig, Metro Manila, Philippines	PGH, Metro Manila Tel. No. (02)524 1078 (02) 554 8400 loc 2311	B. Rodriguez St., Cebu City Tel. No. (032) 253 9891 loc 353	Bajada, Davao City Tel. No. (082) 227 2731
Tel. No. (02) 811 8000 Fax No. (02) 838 1025 Emergency Hotline (02) 831 5576			



BASF
We create chemistry

Kotetsu®



For the effective control of Diamondback Moth, Aphids, Cabbage Moth, Thrips and Flower Thrips (Banana)

Compositions

Chlorfenapyr 100 g/L
Water 900 ml/L

Net Content:
1 Liter

Registered with the Fertilizer and Pesticide Authority Pursuant to P.D. 1144
FPA Registration Number: 005-283-3763

DIRECTIONS FOR USE

Crop /Pests	Tbsp / 16L	Application
CABBAGE Diamond back moth Aphids Cabbage moth	1.5 – 3.0	Start spraying during early stage of infestation. Repeat at 7 days interval or as needed.
POTATO Thrips	0.5 – 1.5	Start spraying during early stage of infestation. Repeat at 7 to 14 days interval or as needed.
BANANA Flower thrips	2-3 mL/ 1L water	Flower bud injection. Inject 60-80 mL solution per bud when at least ¼ inflorescence has emerged.

1 tbsp= 10ml

RE-ENTRY PERIOD: Do not enter the field within 2 days after spraying.

PRE-HARVEST INTERVAL: Do not spray 14 days before harvest

COMPATIBILITY: KOTETSU 10SC can be mixed with most commonly used insecticides and fungicides.

DIRECTION FOR STORAGE AND DISPOSAL: Store in original container tightly closed and away from foods and feeds. Keep in a cool and dry place. Do not re-use empty container. Destroy empty container and bury away from water sources.

WARRANTY AND PROHIBITION: It is a violation of FPA rules and regulations to use this product in manner inconsistent with its label. No warranty of any kind, expressed or implied is made concerning the use of this product. Users assumes all risks and liability resulting from handling, use or application.



CAUTION