

 **BASF**
The Chemical Company

Regent[®]
0.3GR

3 g/kg Fipronil

**A Systemic Granular Insecticide for
the control of Stemborers and Brown
Planthoppers in Rice; Asiatic Corn
Borer, Earworm and Aphids in Corn;
and White Grubs in Sugarcane**

Registered with the Fertilizer and Pesticide Authority
Pursuant to P.D. 1144
FPA Registration Number: 005-244-0541
® = Registered Trademark of BASF

5 kg

KEEP OUT OF REACH OF CHILDREN



GENERAL INFORMATION: REGENT 0.3GR is a systemic pyrazole insecticide that provides excellent control of stemborers and brown planthoppers in rice; asiatic corn borer, earworm and aphids in corn; and white grubs in sugarcane. It is very effective in controlling brown planthoppers with known resistance to other insecticides. **REGENT 0.3GR** works by contact and stomach action with long residual activity.

DIRECTIONS FOR USE:

CROP	INSECT PESTS	DOSE RATE (KG PROD/HA)	APPLICATION METHOD AND TIMING
Paddy Rice (Transplanted)	Stemborers	10-17	Broadcast evenly at 30 to 45 days after transplanting (DAT)
(Direct Seeded)	Brown Planthoppers	10-17	Broadcast evenly at 40 to 45 days after sowing (DAS)
Corn	Asiatic Corn Borer, Earworm, Aphids	10-13	Apply into the whorl at 30 to 35 days after emergence (DAE)
Sugarcane (Plant canes)	White Grubs	20-30	Apply as infurrow at 50 days after planting (DAP)
(Ratoon)			Apply as infurrow along with the first application of the fertilizer (i.e. after stubble shaving and replanting)

For rice, irrigate the field to a depth of 3 to 5 cm. Close all levees then broadcast **REGENT 0.3GR** evenly.

COMPATIBILITY: Compatible with most types of fertilizers and pesticides.

PRE-HARVEST INTERVAL: Do not apply 7 days before harvest.

RE-ENTRY: One day after application.

STORAGE AND DISPOSAL: Store in original container, tightly closed in a safe place and away from children's reach. Do not store near seeds, food and feedstuffs. Do not store near heat or open flame. Burn or bury empty containers in an isolated area at least 0.5 meter depth and away from sources of water. Do not re-use empty containers.

PROHIBITION AND WARRANTY: It is a violation of FPA rules and regulations to use this product in a manner inconsistent with its label. No warranty of any kind, expressed or implied, is made concerning the use of this product. The consumer assumes all risks and liabilities resulting from the handling, use or application of the product.

PRECAUTIONS: REGENT 0.3GR may irritate eyes. Avoid contact with eyes, skin and clothing. WEAR gloves, dust mask **WHEN HANDLING OR DURING APPLICATION**. Do not eat, drink or smoke when using insecticide. Wash exposed skin thoroughly with soap and water after handling and application. Bath thoroughly after working with this insecticide. Remove and wash thoroughly all protective equipment and working clothes before re-use.

BABALA: Magbibigay ng bahagyang kati sa mata. Iwasan ang matagalang pagkadikit sa balat. Gumamit ng guwantes, respirator at pantrabahong damit na mahaba ang manggas habang nagsasabog ng **REGENT 0.3GR**. Huwag kumain, uminom at manigarilyo habang ginagamit ang produkto. Hugasang mabuti ang mga kamay, braso at mukha. Labhang mabuti ang mga pantrabahong damit at guwantes bago muling gamitin.

SIGNS AND SYMPTOMS OF POISONING: General symptoms of overexposure include respiratory difficulty, lethargy, tremor and in severe cases, convulsion.

FIRST AID TREATMENT: In case one or more of the above symptoms appear, stop working immediately and perform first aid treatment, accordingly:

IF ON SKIN: Remove contaminated clothing. Immediately flush contaminated skin with plenty of water.

IF ON EYES: Flush eyes with steady running water for at least 20 minutes. Get medical attention immediately.

IF INHALED: Remove victim from area of exposure and bring to fresh air. Apply artificial respiration when necessary.

IF SWALLOWED: If patient is conscious, give 2 to 3 glasses of water. Induce vomiting by touching the back of the throat with finger. May give charcoal slurry if available. If patient is unconscious, do not induce vomiting nor give anything by mouth. Get **MEDICAL ATTENTION IMMEDIATELY**.

NOTE TO PHYSICIAN: Treat symptomatically. Fipronil is a nerve GABA inhibitor. No known antidote. Give symptomatic and supportive treatment.

IN CASE OF EMERGENCY, PLEASE CONTACT:	
BASF Philippines, Inc. 103 Progress Avenue, Phase 1 GIZ Carmelray Industrial Park 1 4037 Canlubang, Calamba Laguna, Philippines Tel. No.: (049) 549-0001 / (02) 889-4321	VICENTE SOTTO MEDICAL CENTER B. Rodriguez St., Cebu City Tel. No.: (032) 253-9891
NATIONAL POISON CONTROL AND INFORMATION SERVICE PGH, Manila Tel. No.: (02) 526-4248 / (02) 521-8251	DAVAO MEDICAL CENTER Bajada, Davao City Tel. No.: (082) 227-2731

Batch No.:

Formulation Date:

BASF Philippines, Inc.
103 Progress Avenue, Phase 1 GIZ
Carmelray Industrial Park 1
4037 Canlubang, Calamba
Laguna, Philippines

