



Storm® Blocks 4g

Rodenticide Wax Block Bait

Composition:
Flocoumafen 0.05 g/Kg
Bitrex 0.01 g/Kg
Carriers to 1 Kg



Registered with Fertilizer and Pesticide Authority

FPA Registration No:
Net Content: **10 x 15g**



We create chemistry

PRECAUTIONS: When handling or applying baits, do not eat, drink nor smoke. Wear rubber gloves when handling the product. Keep away from food and drinks. Store baits in its original container tightly closed, in a safe place under lock and key. Keep away from children. Do not place bait on or near exposed food or food utensils, food preparation area or food storage area. Do not use where water could be contaminated. Wash hands after handling and applying baits.

BABALA: Huwag kumain, uminom o manigarilyo habang gumagamit nito. Magsuot ng guwantes. Ilayo sa pagkain at inumin. Itago sa orihinal na lalagyan sa lugar na hindi abot ng mga bata. Huwag gamitin sa lugar na maaring mahawaa ang pagkain at tubig. Maghugas ng kamay matapos gumamit nito.

SIGNS AND SYMPTOMS OF POISONING: Symptoms of poisoning include increased tendency to bleed, bruising, nose or gum bleeds. In severe cases, massive bleeding from internal organs may occur.

FIRST AID: If swallowed, obtain medical attention immediately. If not readily available, induce vomiting by putting a finger down the throat. If on skin, removed contaminated clothing. Wash skin with water, using soap if available. If on eyes, flush affected eye with water..

NOTE TO PHYSICIAN: Vitamin K1 is the antidote. In cases of suspected poisoning, determine the problem in time not less than 18 hours after indigestion. If elevation has occurred, administer Vitamin K1 intravenously in divided doses, 40 mg for adults, and 20 mg for children, daily.

STORAGE AND DISPOSAL: Store in original container, tightly closed, away from foodstuffs and animal feeds and out of reach of children. Dispose unwanted baits and dead rodents by burying. Do not re-use empty containers.



Storm® Blocks 4g

Rodenticide Wax Block Bait

Composition:
Flocoumafen 0.05 g/Kg
Bitrex 0.01 g/Kg
Carriers to 1 Kg



Registered with Fertilizer and Pesticide Authority

FPA Registration No:
Net Content: **10 x 15g**




We create chemistry

DIRECTIONS FOR USE: Bait Placement and Inspection. In applications for rat control, select bait points, such as active rat holes, along runs or walkways, around the perimeter of the field, and under rubbish areas. Establish a line of bait points between the rats living and feeding areas. Bait points maybe as close as 5 meters apart in area of heavy infestation. Secure baits at each location on wire or nails, or inside Rodent bait boxes. Mice usually ranges over very limited areas and do not need to drink. They are more difficult to control than rats because their feeding patterns are more random. For control of mice, place baits in areas where activity is obvious, particularly where droppings are seen, One Storm is applied, the bait points should be examined within 7 days, and any baits which have been eaten must be replaced. Baits which have not been eaten must be repositioned. Insdpection of all bait points and rat holes should be repeated at 7-day intervals, placing fresh blocks on each occassion only when blocks have been eaten. **Application Rate:** To control rats and mice in rice, corn, sugarcane, pineapple, banana and other crops, place 2-3 blocks at 5meter interval.

NOTICE TO BUYER: No warranty of any kind, expressed or implied, is made concerning the use of this product. User assumes all risks and liabilities resulting from handling, use or application.

IN CASE OF EMERGENCY, PLEASE CONTACT:			
BASF Philippines, Inc. 11/F HMC Building, 1128 University Parkway North, Ortigas Global City, 1634, Taguig, Metro Manila Tel. (02) 811 8000; Fax: (02) 838 1025; Emergency Contact No.: (02) 831 5578		National Poison Management and Control Center PC-1, Manila Tel.: (02) 524 1078; (02) 554 8400 loc 2311	
Eastern Visayas Regional Medical Center Magsaysay Boulevard, Tacloban City, Leyte Tel.: (053) 321 3131		Zamboanga City Medical Center Dr. Evangelista St., Sta. Catalina, Zamboanga City Tel.: (092) 891 2445; (092) 891 0573	



110 mm

95 mm

130 mm